

Photonic Probes of the Deuteron in Effective Field Theory^a

Silas R. Beane

*Department of Physics, University of Washington,
Seattle, WA 98195.*

I give a taste of recent progress in using chiral perturbation theory to understand nucleon and deuteron structure.

A great deal of progress has been made unravelling nuclear scales and constructing the effective field theory (EFT) relevant for low-energy nuclear physics (for a recent review see Ref. ¹). Here I will focus on a success and a puzzle.

Neutral pion photoproduction on the deuteron has been computed to $O(q^4)$ in chiral perturbation theory (χ_{PT})². The *prediction* for the threshold amplitude is $E_d = (-1.8 \pm 0.2) \times 10^{-3}/m_{\pi^+}$. To see the sensitivity to the elementary neutron amplitude, we set it to zero and find $E_d = -2.6 \times 10^{-3}/m_{\pi^+}$, which results in a cross-section twice as large as the χ_{PT} prediction. This is a unique situation as the EFT makes a prediction that differs significantly from conventional models. A measurement of the threshold amplitude has recently been carried out at Saskatoon³. The experimental results for the pion photoproduction cross-section near threshold are shown in Fig. 1, together with the χ_{PT} prediction at threshold. At threshold, Ref. ³ extracts $E_d = (-1.45 \pm 0.09) \times 10^{-3}/m_{\pi^+}$. While agreement with χ_{PT} to order $O(q^4)$ is not better than a reasonable estimate of higher-order terms, it is clearly superior to models. This is compelling evidence of the importance of chiral loops and of the consistency of nuclear EFT.

Electroproduction on the deuteron has recently been measured at Mainz⁴. The two relevant multipoles have been computed to $O(q^3)$ in the chiral expansion⁵. There are discrepancies between data and theory. However, a true test awaits an $O(q^4)$ computation which is currently underway.

Compton scattering on the deuteron is important because it may provide the only rigorous way to determine the polarizabilities of the neutron. This process has been computed to $O(q^3)$ in χ_{PT} ⁶. An $O(q^4)$ computation is currently underway. The $O(q^3)$ result is in agreement with the 69 MeV data (see Fig. 2). There are $O(q^4)$ counterterms which shift the nucleon polarizabilities. We use this freedom to fit the 95 MeV data using an incomplete $O(q^4)$ calculation. A reasonable fit at backward angles can be achieved with $\alpha_n = 4.4$ and

^aTalk given at the Workshop on Chiral Dynamics 2000: Theory and Experiment, TJNL (Newport News, USA) 17th – 22nd July 2000; to be published in the Proceedings. Preprint NT@UW-00-032.




Figure 1: Total cross section for $\gamma d \rightarrow \pi^0 d$. The data from SAL are depicted by the boxes, the CHPT threshold prediction (before the data release) is the star on the dotted line (indicating the threshold photon energy).

$\beta_n = 10$, which are in startling disagreement with the $O(q^3)$ χ_{PT} expectations. Also, the cross-section at 69 MeV with $\alpha_n = 4.4$ and $\beta_n = 10$ is then excessively forward peaked and misses the data. This situation poses an interesting puzzle. Of course, a full $O(q^4)$ calculation in χ_{PT} is necessary if the neutron polarizability is to be extracted from the Saskatoon data in a systematic way.




Figure 2: The differential cross section for elastic γ -deuteron Compton scattering at incident photon energies of $E_\gamma = 69$ MeV and 95 MeV compared to data from Ref. ⁷ and Ref. ⁸, respectively.

1. S.R. Beane *et. al.*, nucl-th/0008064.
2. S.R. Beane *et. al.*, Nucl. Phys. **A618**, 381 (1997).
3. J.C. Bergstrom *et. al.*, Phys. Rev. **C57**, 3203 (1998).
4. I. Ewald *et. al.*, nucl-ex/0010008
5. V. Bernard, H. Krebs, Ulf-G. Meißner, Phys. Rev. **C61**, 058201 (2000).
6. S.R. Beane *et. al.*, Nucl. Phys. **A656**, 367 (1999).
7. M. Lucas, Ph. D. thesis, University of Illinois, unpublished (1994).
8. D.L. Hornidge *et. al.*, Phys. Rev. Lett. **84**, 2334 (2000).